

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0311F

October 27, 1983

TESTIMONY BY

Samuel Milham, Jr., M.D., M.P.H.

REPRESENTING

Washington State Department of
Social and Health Services

BEFORE

Environmental Protection Agency Hearing on
Proposed Standards for High-Arsenic Copper Smelters
November 2, 1983

My name is Samuel Milham, Jr. I am a physician employed as Head of the Epidemiology Section at the Washington State Department of Social and Health Services.

In the early 1970s, children living near a lead smelter in Texas were shown to have high blood lead levels. In 1972, DSHS launched a series of studies to investigate the nature of exposure to heavy metals in people living in Ruston, near the ASARCO Copper Smelter.

Initial studies indicated that blood lead levels and blood enzymes affected by lead were within normal limits. However, arsenic levels in urine and hair in children residing near the Smelter were elevated as compared to children who resided at a distance (8 miles) from the Smelter¹. Levels of arsenic in urine, house dust, and soil were found to decrease rapidly with distance of residence from the Smelter. Urinary arsenic levels varied synchronously over a 5-week period suggesting that inhalation was the most likely route of exposure. A decreasing linear relationship is seen between urinary arsenic and increasing age with younger children having consistently higher levels.

AR 18.3.8 0006

-2-

During a strike at the Tacoma Smelter in 1974, urinary arsenic levels in the community were lower than when processing resumed, suggesting a direct impact of the Smelter arsenic emissions on human exposures, most likely mediated by inhalation². Urinary arsenic levels in the years since 1972 have shown variation over time, but no clear time trend is demonstrated in the urinary arsenic levels of Ruston children. In 1975, Ruston children averaged 35 micrograms of arsenic per liter of urine; in 1983, they averaged 36 micrograms of arsenic per liter of urine.

In an attempt to assess the health impact of arsenic exposure in the community around the ASARCO Tacoma Smelter, a number of studies have been done:

1. Absenteeism in Ruston Elementary School was found to be no different than in 6 other Tacoma elementary schools³.
2. Pure tone hearing screening tests done in the Ruston Elementary School gave similar results to those done at other Tacoma elementary schools³.
3. Pure tone threshold audiometry done on 7 Ruston children with high urinary arsenic levels ($\geq .2$ PPM on 2 or more sample days) was normal³.
4. Average blood values of 33 Ruston Elementary School children were found to be the same as those of 25 control children (Fern Hill Elementary School).
5. Chromosome analysis (sister chromatid exchange) was normal in 5 arsenic exposed Ruston children and in 5 unexposed controls.
6. Growth and development of Ruston School children, as measured by height and weight attained at a given age, was found to agree with U.S. averages. Academic and physical performance of Ruston Elementary School children was similar to that of other Tacoma elementary school children.
7. Mortality due to lung cancer in the census tracts near the Smelter was not elevated compared to more distant tracts for deaths in the years 1950-1970.
8. Follow up of children enrolled at Ruston Elementary School for 3 or more years during the years 1900-1919 was attempted. Survivorship of 137 males in the group was found to be favorable (more survivors to 1980 than expected).

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

03 13

-3-

Published studies of health effects in the communities around other arsenic emitters⁴⁻¹³ are all essentially negative.

In spite of the failure to date to delineate any adverse health effects due to arsenic in the community around the ASARCO Tacoma Smelter, I feel that it would be prudent to minimize human exposure to arsenic by reducing arsenical emissions to the lowest level possible. It is especially important that low-level or fugitive emissions be reduced. To this end, I recommend:

1. Setting a community 24-hour ambient air arsenic standard.
2. Establishing an air sampling network in the impacted communities to monitor ambient air arsenic.
3. Monitoring urinary arsenic levels of people residing in the impacted communities on a regular basis.

-4-

References

1. Milham, S. Jr. and Strong, T.R. Human Arsenic Exposure in Relation to a Copper Smelter. Environmental Research 7: 176-182, 1974.
2. Milham, S. Jr. Industrial Toxins and the Community. Arsenic Contamination Near a Copper Smelter. Occupational Safety and Health Symposia 1977. DHEW (NIOSH) Publication No. 78-169, 1978.
3. Milham, S. Jr. Studies of Morbidity Near a Copper Smelter. Environmental Health Perspectives 19: 131-132, 1977.
4. Blott, W.J. and Fraumeni, J.F. Jr. Arsenical air pollution and lung cancer. The Lancet, July 26, 1975, 142-144.
5. Lyon, J.L., Fillmore, J.L. and Klauber, M.R. Arsenical air pollution and lung cancer. The Lancet, October 22, 1977, 869.
6. Rom, W.N., Varley, G., Lyon, J.L. and Shopkow, S. Lung Cancer Mortality among residents living near the El Paso Smelter. British Journal of Industrial Medicine 39, 269-272, 1983.
7. Greaves, W.W., Rom, W.N., Lyon, J.L., Varley, G., Wright, D.D., and Chiu, G. Relationship between lung cancer and distance of residence from nonferrous smelter stack effluent. American Journal of Industrial Medicine 2, 15-23, 1981.
8. Pershagen, G., Elinder, C.G., and Bolander, A.M. Mortality in a region surrounding an arsenic emitting plant. Environmental Health Perspectives 19, 133-137, August 1977.
9. Matanoski, G., Landau, E., Tonascia, J., Lazar, C., Elliot, E., McEnroe, W. and King, K. Cancer mortality in an industrial area of Baltimore. Environmental Research 25, 8-28, 1981.
10. Morse, D.L., Harrington, J.M., Housworth, J., and Landrigan, P.J. Arsenic exposure in multiple environmental media in children near a smelter. Clinical Toxicology 14(4), 389-399, 1979.
11. Baker, E.L., Hayes, C.G., Landrigan, P.J., Handke, J.L., Leger, R.T., Housworth, W.J. and Harrington, J.M. A nationwide survey of heavy metal absorption in children living near primary copper, lead and zinc smelters. American Journal of Epidemiology 106(4), 261-273, 1977.

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

03 15

-5-


12. Nordstrom, S., Beckman, L. and Nordenson, I. Occupational and Environmental risks in and around a smelter in northern Sweden, VI. Congenital Malformations. Hereditas 90, 297-302, 1979.
13. Nordstrom, S., Beckman, L. and Nordenson, I. Occupational and environmental risks in and around a smelter in northern Sweden, V. Spontaneous Abortion among female employees and decreased birth weight in their offspring. Hereditas 90, 291-296, 1979.

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0316




Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0317

FIGURE 1 Location of Tacoma Smelter and Ruston In Relation to the City of Tacoma



Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0318

BLOOD LEAD BY SCHOOL
JUNE 7, 1972

| UG/100 ML | NUMBER OF STUDENTS | |
|-----------|--------------------|----------------|
| | RUSTON | FERN HILL |
| 0-4 | 0 | 0 |
| 5-9 | 12 | 5 |
| 10-14 | 2 | 6 |
| 15-19 | 9 | 7 |
| 20-24 | 2 | 3 |
| 25-29 | 2 | 1 |
| 30+ | 0 | 0 |
| TOTAL | 27 | 22 |
| MEAN | 14.7 ug/100 ML | 15.8 ug/100 ML |


Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0317

Fig.2 URINARY ARSENIC BY SCHOOL



ASA219





Fig. 5 URINARY ARSENIC IN RUSTON CHILDREN FOR 5 WEEKLY SAMPLES




Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0321

Fig.4 URINARY ARSENIC BY DISTANCE FROM SMELTER




0322

ASA219

ADMINISTRATIVE REMOVAL

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.



Notice: If the film image
is less clear than this
notice, it is due to the
quality of the document
being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0323

TABLE 1

Urinary and House Dust Arsenic by Distance from Smelter

| <u>Distance of Residence from Stack</u> | <u>Mean Urinary Arsenic in PPM</u> | <u>Vacuum Cleaner dust Arsenic in PPM</u> | <u>Attic Dust Arsenic in PPM</u> |
|---|--|---|--------------------------------------|
| 0 - .4 miles | .30 | 1300 | 2100 |
| .5 - .9 miles | .19 | 970 | |
| 1.0 - 1.4 miles | .08 | 330 | |
| 1.5 - 2.0 miles | .06 | no sample | |
| 2.0 - 2.4 miles | .02 | 70 | |

e: Prevailing wind from the north at this sampling

Notice: If the film image
is less clear than this
notice, it is due to the
quality of the document
being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0324

Figure 1 is a line graph showing the relationship between P.P.M. ARSENIC (Y-axis) and AGE (X-axis). The Y-axis ranges from 0 to 0.20, and the X-axis ranges from 2 to 12. The graph illustrates arsenic levels in siblings, with lines connecting siblings. Data points are marked with various symbols (circles, squares, triangles) and connected by lines. The graph shows a general downward trend in arsenic levels as age increases, with some variability between siblings.

| Age | P.P.M. ARSENIC | Symbol |
|-----|----------------|----------|
| 4 | 0.19 | Circle |
| 6 | 0.15 | Circle |
| 7 | 0.11 | Circle |
| 8 | 0.19 | Square |
| 10 | 0.08 | Square |
| 10 | 0.08 | Circle |
| 10 | 0.07 | Square |
| 10 | 0.04 | Triangle |
| 10 | 0.09 | Square |
| 10 | 0.08 | Circle |
| 3 | 0.13 | Triangle |
| 8 | 0.09 | Triangle |

52E0

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.


ADMINISTRATIVE REMOVAL

ASA219

0326

FIGURE 1

TOTAL URINARY ARSENIC BY AGE



Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0327

AVERAGE URINARY ARSENIC LEVELS (PPM)
DURING AND AFTER STRIKE AT THE
TACOMA SMELTER 1974

| | DURING | | | AFTER | | |
|----------|----------------|----------------|----------------|----------------|----------------|----------------|
| | <u>7-19-74</u> | <u>7-25-74</u> | <u>AVERAGE</u> | <u>8-20-74</u> | <u>8-22-74</u> | <u>AVERAGE</u> |
| CHILD 1 | .17 | .34 | .25 | .68 | .63 | .655 |
| CHILD 2 | .27 | .07 | .17 | .16 | .41 | .285 |
| CHILD 3 | .01 | .08 | .045 | .11 | .06 | .085 |
| CHILD 4 | x | .14 | .14 | .10 | .13 | .115 |
| CHILD 5 | .01 | .04 | .025 | .03 | .05 | .04 |
| CHILD 6 | .01 | .04 | .025 | .04 | .08 | .06 |
| CHILD 7 | .04 | .08 | .06 | .09 | .10 | .095 |
| CHILD 8 | .03 | .08 | .055 | .06 | .05 | .055 |
| CHILD 9 | x | .06 | .06 | x | .22 | .22 |
| CHILD 10 | .02 | .01 | .015 | .02 | .03 | .025 |
| ALL | | | .08 | | | .16 |

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.


ADMINISTRATIVE REMOVAL

ASA219

0328

FIGURE 2

Correlation of As^{+3} and DMAA in individual urine specimens



Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0327

Chemical Forms of Arsenic in Urine

Average concentration for 23 Ruston residents by age

| <u>Age.</u> | <u>N</u> | <u>As⁺³</u> | <u>As⁺⁵</u> | <u>MAA</u> | <u>DMAA</u> |
|-------------|----------|------------------------|------------------------|------------|-------------|
| 2-7 | 6 | 4.5 | 2.9 | 5.5 | 63.1 |
| 8-9 | 5 | 3.5 | 1.1 | 3.2 | 28.6 |
| 10-14 | 4 | 2.9 | .3 | 3.1 | 26.5 |
| Adult | 8 | 1.0 | .4 | .7 | 12.0* |

* 4.2 without 1 outlying result

MAA = methylarsonic acid

DMAA = dimethylarsinic acid

TABLE 2
URINARY ARSENIC (PARTS PER MILLION) FOR RUSTON PRESCHOOL CHILDREN
1976

| Age | Date/Day | | | | | | | Average |
|---------|---------------|--------------|--------------|--------------|--------------|---------------|-------------|---------|
| | 8/26 Thur. | 8/27 Fri. | 8/28 Sat. | 8/29 Sun. | 8/30 Mon. | 8/31 Tues. | 9/1 Wed. | |
| 4 | .06 | .05 | .16 | .09 | .22 | .03 | .03 | .091 |
| 4 | .10 | .14 | .12 | .10 | .30 | .12 | .10 | .140 |
| 5 | .09 | .05 | .14 | .10 | X | .04 | .06 | .080 |
| 4 | .02 | .02 | .01 | .02 | .02 | .03 | .03 | .021 |
| 5 | .40 | .40 | .15 | .27 SF | .25 | .31 | .16 | .277 |
| 3 | .10 | .01 | .08 | .11 SF | X | .03 | .36 | .115 |
| Average | .128 | .112 | .110 | .115 | .198 | .093 | .123 | .122 |

X = no specimen
SF = ate shrimp

0000

ASA219

ADMINISTRATIVE REMOVAL

Notice: If the film image
is less clear than this
notice, it is due to the
quality of the document
being filmed.

TABLE 3
URINARY ARSENIC (PARTS PER MILLION) FOR RUSTON SCHOOL CHILDREN
1976

| | Age | Date/Day | | | | | | | | | | Average |
|--------------|-----|--------------|--------------|--------------|--------------|---------------|--------------|---------------|--------------|--------------|-------------|---------|
| | | 7/23 Fri. | 7/24 Sat. | 7/25 Sun. | 7/26 Mon. | 7/27 Tues. | 7/28 Wed. | 7/29 Thur. | 7/30 Fri. | 7/31 Sat. | 8/1 Sun. | |
| Family No. 1 | | | | | | | | | | | | |
| Child 1 | 10 | .03 | .03 | .02 | .05 | .02 | .08 | .08 | .03 | X | X | .043 |
| Child 2 | 9 | .03 | .08 | .04 | .06 | .07 | .04 | .04 | X | X | X | .051 |
| Child 3 | 12 | .03 | .04 | .04 | .04 | .03 | .03 | .03 | .01 | X | X | .031 |
| Family No. 2 | | | | | | | | | | | | |
| Child 1 | 8 | .16 | .11 | .05 | .08 | .13 | .26 | .11 | .08 | .10 | .07 | .115 |
| Child 2 | 10 | .04 | .08 | .04 | .07 | .07 | .06 | .09 | .04 | .05 | .11 | .065 |
| Child 3 | 6 | .14 | .22 | .10 | .17 | .12 | .22 | .07 | .06 | .19 | .07 | .136 |
| Family No. 3 | | | | | | | | | | | | |
| Child 1 | 6 | .05 | .11 | .03 SF | .59 | .13 | .04 | .18 | .15 | .20 | .80 | .228 |
| Child 2 | 9 | .09 | .10 | .09 | .06 | .07 | .18 | .17 | .06 | .17 | .19 | .118 |
| Child 3 | 10 | .04 | .10 | .03 | .05 | .10 | .08 | .06 | .07 | .04 | .46 | .103 |
| Child 4 | 7 | .10 | .09 | .06 SF | .20 | .06 | .11 | .14 | .17 | .20 | .89 | .202 |
| Average | | .071 | .096 | .050 | .135 | .080 | .112 | .097 | .074 | .136 | .370 | .114 |

SF = ate salmon
(= no specimen

ASA219

ADMINISTRATIVE REMOVAL

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

03332

| Sampling Date | Number of individuals sampled | Group Studied | Minimum | Maximum | Average |
|------------------------|-------------------------------|---|---------|---------|---------|
| 6-6-72 | 19 | Ruston School | 10 | 150 | 81.8 |
| 6-7-72 | 16 | Fern Hill School | 10 | 50 | 20.0 |
| 6-7-72 | 9 | Ruston Preschool Children | 40 | 620 | 270.0 |
| 9-12-72 | | Traverse Study of Ruston and Tacoma within: | | | |
| | 7 | .5 mi. of stack | 50 | 620 | 300.0 |
| | 8 | .5 - 1.0 mi. of stack | 50 | 420 | 190.0 |
| | 6 | 1.0 - 1.5 mi. of stack | 40 | 140 | 80.0 |
| | 5 | 1.5 - 2.0 mi. of stack | 40 | 100 | 60.0 |
| | 6 | 2.0 - 2.5 mi. of stack | N.D. | 50 | 20.0 |
| | 5 | 2.5 - 3.0 mi. of stack | 10 | 100 | 46.0 |
| | 5 | 3.0 - 3.5 mi. of stack | 10 | 50 | 34.0 |
| | 10 | 3.5 - 4.0 mi. of stack | 10 | 110 | 48.0 |
| 10-11-72 | | Ruston Children | | | |
| 10-25-72 | | Average of 5 weekly samples | | | |
| 10-30-72 | 14 | | 20 | 470 | 99.0 |
| 11-6-72 | | | | | |
| 11-13-72 | | | | | |
| 9-18-73 | 107 | Ruston School Children | N.D. | 430 | 81.0 |
| 10-25-73 | 106 | Ruston School Children | 10 | 470 | 55.0 |
| 7-19-74 | 8 | Ruston Children (Smelter on strike) | 10 | 270 | 70.0 |
| 7-25-74 | 10 | | 10 | 340 | 94.0 |
| 8-20-74 | 9 | Ruston Children (after Smelter strike) | 20 | 680 | 145.0 |
| 8-22-74 | 10 | | 30 | 630 | 176.0 |
| 6-3-75 | | Ruston School Children | | | |
| | 5 | Seafood ingestion | 30 | 190 | 102.0 |
| | 36 | No seafood ingestion | 20 | 660 | 87.0 |
| 6-3-75 | | Fern Hill School Children | | | |
| | 13 | Seafood ingestion | 10 | 270 | 62.0 |
| | 48 | No seafood ingestion | 10 | 230 | 25.0 |
| 11-17-75 | 102 | Ruston School Children | 10 | 200 | 40.0 |
| | 17 | Seafood ingestion | 10 | 150 | 68.0 |
| | 85 | No seafood ingestion | 10 | 200 | 35.0 |
| 7-23-76 through 8-1-76 | 10 | Ruston School Children for 10 days each | 20 | 890 | 114.0 |
| 8-26-76 through 9-1-76 | 6 | Ruston Preschool children for 7 days each | 10 | 400 | 122.0 |
| 6-30-83 | 22 | N. Tacoma Children | 10 | 116 | 36.0 |
| | 27 | Vashon Island Children | < 10 | 116 | 23.0 |
| | 22 | Olympia Children | < 10 | 87 | 12.0 |

RECOMMENDATIONS

1. SETTING A COMMUNITY 24-HOUR AMBIENT AIR ARSENIC STANDARD
2. ESTABLISHING AN AIR SAMPLING NETWORK IN THE IMPACTED COMMUNITIES
TO MONITOR AMBIENT AIR ARSENIC
3. MONITORING URINARY ARSENIC LEVELS OF PEOPLE RESIDING IN THE
IMPACTED COMMUNITIES ON A REGULAR BASIS

0333

ASA219

ADMINISTRATIVE REMOVAL

Notice: If the film image
is less clear than this
notice, it is due to the
quality of the document
being filmed.

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0334

Table 1

1969-74 Average Annual

| <u>School</u> | <u>Attendance (A)</u> | <u>Enrollment (B)</u> | <u>A/B</u> |
|----------------|-----------------------|-----------------------|------------|
| Ruston | 134 | 141 | .95 |
| Sherman | 591 | 637 | .93 |
| Point Defiance | 451 | 477 | .95 |
| Truman | 578 | 605 | .96 |
| Fern Hill | 650 | 688 | .94 |
| Larchmont | 342 | 362 | .94 |
| Oakland | 189 | 201 | .94 |

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0335

MORBIDITY STUDIES

PURE TONE HEARING SCREENING TESTS DONE IN THE RUSTON ELEMENTARY SCHOOL GAVE SIMILAR RESULTS TO THOSE DONE AT OTHER TACOMA ELEMENTARY SCHOOLS.

PURE TONE THRESHOLD AUDIOMETRY DONE ON 7 RUSTON CHILDREN WITH HIGH URINARY ARSENIC LEVELS ($\geq .2$ PPM ON 2 OR MORE SAMPLE DAYS) WAS NORMAL.

CHROMOSOME ANALYSIS (SISTER CHROMATID EXCHANGE) WAS NORMAL IN 5 ARSENIC EXPOSED RUSTON CHILDREN AND IN 5 UNEXPOSED CONTROLS.

GROWTH AND DEVELOPMENT OF RUSTON SCHOOL CHILDREN, AS MEASURED BY HEIGHT AND WEIGHT ATTAINED AT A GIVEN AGE, WAS FOUND TO AGREE WITH U.S. AVERAGES. ACADEMIC AND PHYSICAL PERFORMANCE OF RUSTON ELEMENTARY SCHOOL CHILDREN WAS SIMILAR TO THAT OF OTHER TACOMA ELEMENTARY SCHOOL CHILDREN.

Notice: If the film image is less clear than this notice, it is due to the quality of the document being filmed.

ADMINISTRATIVE REMOVAL

ASA219

0336

Table 3

Average Blood Values, Ruston and Fern Hill Schools, Grades 3-5
May, 1975

| | Ruston | Fern Hill | Normal Values (Wintrobe(4)). | |
|--|--------|-----------|---------------------------------|-----------|
| | | | Age 6-10 | Age 11-15 |
| Red Blood Count(millions/cu.mm.) | 4.48 | 4.71 | 4.7 | 4.8 |
| Hemoglobin(gm/100 ml) | 13.0 | 13.6 | 12.9 | 13.4 |
| Hematocrit(vol. of packed RBC/100 ml) | 37.2 | 38.8 | 37.5 | 39.0 |
| Mean Corpuscular Vol.(cu) | 82.4 | 81.8 | 80 | 82 |
| Mean Corpuscular Hemoglobin(γγ) | 28.9 | 29.9 | 27. | 28 |
| Mean Corpuscular Hemoglobin concentration(%) | 35.3 | 36.7 | 34 | 34 |
| White blood count (per cu. mm.) | 5,720 | 6,080 | 8,100 | 8,000 |
| Percent granulocytes | 46.3 | 49.0 | 50 | 51 |
| Percent lymphocytes | 47.6 | 43.6 | 39 | 38 |


ADMINISTRATIVE REMOVAL

ASA219

03371

Lung Cancer Mortality Near a Copper Smelter - Hartley

Figure 1
City of Tacoma and Ruston



Significant ($P < .05$) SMR elevations for lung cancer